

# Al-Razi University Journal for Medical Sciences



## RUJMS

ISSN No. 2616-6143

Volume (3) Issue (1) June 2019



مرخصة من وزارة التعليم العالي والبحث العلمي  
بقرار وزاري رقم ( 57 ) لسنة 2018م

RUJMS

Published by Al-Razi University

Bianual Referred Journal

All Rights Reserved for Al-Razi University

---

Editor in Chief	Nationality	Degree
Dr. Nabil Ahmed Al-Rabeei	Yemen	Professor
Editor Manager	Nationality	Degree
Dr. Rashad Al-Namer	Yemen	Associate Professor

### Editorial Board Members

No	Editorial board members	Nationality	Degree
1.	Dr. Abdulsalam.M. Dallak	Yemen	Professor
2.	Dr. Abduljalil D. Ghaleb	Yemen	Professor
3.	Dr. Mohammed Abdulhaleem	Yemen	Professor
4.	Dr. Mohammed Aissa	Yemen	Professor
5.	Dr. Ahmed Al-Sobati	Yemen	Professor
6.	Dr. Abdulhameed Al-Thifani	Yemen	Associate Professor
7.	Dr. Nouradden Al-Jaber	Yemen	Associate Professor
8.	Dr. Shatha Hassan Yassin	Yemen	Associate Professor
9.	Dr. Sadeq Hassan Al-Sheraji	Yemen	Associate Professor
10.	Dr. Ahmed Ali Abdulateef	Yemen	Associate Professor
11.	Dr. Abdulmajid Alssaifi	Yemen	Associate Professor

### Advisory Board

No	Advisory Board	Nationality	Degree
1.	Dr. El Houcin Boidida	Morocco	Associate Professor
2.	Dr. Yahia Cherrah	Morocco	Associate Professor
3.	Dr. Abdulaziz Benjouad	Morocco	Associate Professor
4.	Dr. Abdellah Akil	USA	Associate Professor
5.	Dr. Katim Alaoui	Morocco	Associate Professor
6.	Dr. Arvinder Bahala	India	Associate Professor
7.	Dr. David Tasala	USA	Associate Professor

**Copyright of articles published in the RUJMS belong to the University of Al-Razi unless the work is subject to copyright.**

**Address: Al-Razi University - College of Medical Sciences**

**Telefax: +9671406760 P.O. Box:1152 Sana'a – Yemen**

**Website: <http://alraziuni.edu.ye/rujms/>**

**Designed by Eng. Osama Al-Moaina  
Ossamah245@yahoo.com**

## TABLE OF CONTENTS

<b>ORIGINAL ARTICLES</b>	<b>PAGE</b>
<b>1. Tumors of the Larynx in Yemen: Prevalence and Treatment</b> Ali Obaid Muthanna	<b>1-7</b>
<b>2. Effect of Training Program on Nurses ' Knowledge Toward Care of Patients With Myocardial Infarction in Al-Thowrah Hospital, Al-Hodeida City, Yemen</b> Sadek Abdu Alwsaby, Nabil Ahmed Al-Rabeei, Abdelaziz Baalawi and Ali Floos	<b>8-15</b>
<b>3. Mothers' Knowledge Toward Malnutrition of Children Under 5 Years Old in Al-Sabeen Hospital, Sana'a City- Yemen</b> Sadek Abdu Alwsaby, Nabil Ahmed Al-Rabeei, Abdelaziz Ahmed Baalawi, and Ali Ahmed Floos	<b>16-22</b>
<b>4. Assessment of Compliance of Written Pharmaceutical Advertisements in Sana'a-Yemen to Criteria of World Health Organization</b> Anes A. M. Thabit, Saleh Yagob, Abdulaziz Mofid, Abdullah Alomaesi, Abdullah Helal, Akram Qahtan, Akram Almasanea, Amar Alsoudi, Bassam Azzam, Mohammed Seraj, Saddam Hashem, Zkee Almacheheri.	<b>23-27</b>
<b>5. Medical Treatment for Orbital Complications Secondary to Rhinosinusitis</b> Ali Obaid Muthanna	<b>28-34</b>
<b>6. Comparative Study of in vitro Quality Specifications of Yemeni Brand of Glimepiride Tablets Versus Foreign Brands Marketed in Yemen</b> Alaa Abdulkarim Almaqtari and Anes A. M. Thabit	<b>35-40</b>



## Medical Treatment for Orbital Complications Secondary to Rhinosinusitis

Ali Obaid Muthanna

Otolaryngology, Head & Neck Surgery Department, Faculty of Medicine and Health Sciences, Sana'a University, Yemen

\*Corresponding author: email: muthannadr@yahoo.com

### Abstract

**Background:** Despite the modern antibiotherapies applied in the practice of otorhinology, the orbital complications of sinusitis are still considered a serious threat to essential functions of the eye, including loss of vision and life threatening symptoms. **Aim.** To assess the efficiency of medical treatment in orbital complications secondary to rhinosinusitis. **Methods:** Quasi-experimental study was admonished among 30 patients with orbital complications secondary to rhinosinusitis from January 2010-June 2017, at OtoRhinoLaryngology, Head & Neck Department, at Al-Thowrah general modern hospital, Sana'a, Yemen. All patients underwent to clinical examination, radiological investigations (C T, MRI), and initially aggressive medical treatment by IV. Antibiotic and locally nasal decongestant for 24-48 hours, if there is no evidence of improvement, we intervened with a surgical procedure, by nasal endoscopy or external approach in the form of external ethmoidectomy to evacuate collection of pus. **Results:** A total of 30 patients had complained of orbital complications secondary to rhinosinusitis were included in the study. Age ranged 2-50 year, males (73.3%) and females (26.7%) Patients. Children were the most category affected 24 patients, right was more than left side ethmoidal sinus was the most sinus affected, followed by ethmoid and maxillary sinuses together. Preseptal cellulitis was the commonest complication (53.3%) patients, orbital cellulitis, and subperiosteal abscess occurred in (20%) patients respectively. Medical treatment successes in (80%) patients, surgical procedure under covering of antibiotic was done in (20%) patients. **Conclusion:** Orbital complications were the most common complications secondary to rhinosinusitis and were more in the children. The outcome of the treatment depended on the types of complications. Medical treatment was showed high efficiency rate.

**Keywords:** Orbital complications, rhinosinusitis, surgical procedure, medical treatment.

### Introduction

Close proximity of the orbit to the paranasal sinuses makes it the most commonly involved structure in complications of rhinosinusitis. Congenital dehiscence in the lamina papyracea which separates the ethmoid sinuses from the orbit exposes the orbital contents to direct extension of rhinosinusitis.<sup>1</sup> In addition, the superior

and inferior ophthalmic veins that are valveless, allow direct communication between the nose, ethmoid sinuses, orbit and cavernous sinus and spread of infection. Orbital involvement primarily results from a thrombophlebitis and interference with the venous drainage of the orbital contents.<sup>2,3</sup>

Children appear to be more prone to orbital complications of rhinosinusitis, probably because of their higher rates of upper respiratory tract infection and rhinosinusitis.<sup>1</sup>

The orbital complications of rhinosinusitis are classified into five categories; preseptal, orbital cellulitis, subperiosteal abscess, orbital abscess, and cavernous sinus thrombosis. Chandler et al (1970)<sup>2</sup>.

The incidence of orbital complications secondary to rhinosinusitis has been reported between 21 and 90% but do not appear to be altered during the last 20 years.<sup>4,5</sup> The methods for the treatment of these complications today often debated: conservative, surgical or combined. Surgical options include both endoscopic and open surgery. However, in the era of endoscopic surgery of the paranasal sinuses, there are still no clear advantage of these technique over open intervention.<sup>6</sup> Some studies reported the use of both technique, while some use either, without any comment on the value of these technique.<sup>7-9</sup>

### **Aim of the study**

Was to assess the efficiency of medical treatment on the orbital complications secondary to rhinosinusitis.

### **Subjects and Methods**

Quasi-experimental study was carried out on 30 patients proved have orbital complications of rhinosinusitis from January 2010 to June 2017, at Otorhinolaryngology Department, Al-Thowrah General Modern Hospital, (AGMH), Sana'a, Yemen. The patients underwent to clinical examination, anterior rhinoscopy, nasal endoscopy after applying local anesthesia (Xylocain spray 10%), and nasal decongestant drops locally. C T scan of the paranasal coronal and axial, hematology study was done, culture and sensitivity test. The patients

admitted to the hospital. Empirical treatment was started by amoxicillin and clavulanic acid by IV route in dose 50mg / kg/day in divided dose every 8 hours for 5 days, then oral same drug for another 10 days (two weeks), dexamethasone in dose 1 mg/ kg / day in divided dose for 5 days then taping for another 5 days and nasal decongestant locally 3 times/day for 5 days. Close observation and clinical evaluation were performed every day. When an improvement in the patient's condition did not occur within 24-48 hours or there is evidence of deterioration of patient's condition, surgical intervention was performed. Six patients were underwent to surgical intervention, 2 patients by intranasal endoscopy approach, and 4 patients by external approach, using the external ethmoidectomy technique, under covering of antibiotic, this allowed a wide opening, good drainage and clear entry to the nasal cavity. Inclusion criteria, all patients with orbital complications, both sex, and all ages, exclusion criteria, fungal sinusitis, nose & paranasal sinuse malignant, congenital anomalies, intracranial complications, uncontrolled diabetes mellitus, and traumatic lesions of the nose.

Ophthalmic consultation was done to assess vision and response to the medical treatment. Data were analyzed using SPSS, Version 20. Frequency and percentage were used.

Chi-square test was used to determine differences between groups, p-value less than 0.05 was considered significant. This study was approved by the ethical committee of our department. Informed consents were obtained from the patients.

### **Results**

A total of 30 patients were enrolled in this study. Males 24 (73.3%) females 8 (26.7%). Age ranged from 2-50 year mean age 11 years .Children (2-12 year) 24 ( 80%), while adults were (30-50 year) 6 (20%) patients. This indicated that the children were the most age categories affected with orbital complications. Right side was more affected than left, 16 to 14 patients respectively. The ethmoid sinuses were most common sinuses affected (80%). Ethmoid and maxillary sinuses were secondary affected. Clinical presentations were showed in


table1. The main presentations were eye lid oedema (Figure 1) pain, and nasal obstruction. Type of complications showed in figure 2. Case of intraorbital abscess has complaining of blindness, while a case with cavernous sinus thrombosis was expired after one week after admission to the hospital. The method of treatment depends on type of complications, 24 patients (80%) underwent to medical treatment and 6 (20%) patients treated by combined modality, medical and surgical. *P*-value less than 0.05. Figure 3.

**Table 1: Clinical presentation of orbital complications among patients (n=30)**


Item	F	%
• Eye lid oedema	30	100
• Pain	30	100
• Nasal obstruction	30	100
• Kemosis	14	47.6
• Tenderness	30	100
• Nasal discharge	20	66.7
• Proptosis	14	47.7
• Ocular movement limitation	14	47.7



**Figure 1: Orbital Edema**



**Figure 2: Type of complications among patients**



**Figure 3: Methods of treatment among patients**

**Discussion**

By virtue of the intimate anatomical relationship between the orbit and the surrounding paranasal sinuses, infection from the sinuses can gain entry into the orbital via several pathways namely through natural defects present in the bony walls that served as partitions between the orbit and paranasal compartments, erosion sinus walls secondary to bony necrosis that resulted from periosteal ischemia induced by the increased intrasinus pressure that occur in acute sinusitis, and via numerous valveless venous channels that provide a further avenue for extension of infection.<sup>8</sup> Despite the

advent of antibiotics and better imaging technique the morbidity of blindness as result of orbital complications has not been fully overcome that is 10% incidence of blindness compared to 20% incidence in pre-antibiotic era.<sup>10</sup> The periosteal is a strong barrier to infection and provides clinicians with time before they are faced with serious complications.<sup>6</sup> Children were the most common category affected in our study, in 80% of patients whereas children have frequented attacks of upper respiratory infections and an immature the immune system. These

results were consistent with previous studies.<sup>4,6,9,11,12</sup>

Male patients' more than female patients without clear explanation for this observation. It is believed that the female immune system is more proficient than that of males.<sup>13</sup>

The ethmoid sinus is most commonly seen to be involved in orbital complications especially in children. This is attributed to close relation between the orbit and the ethmoid sinus with thin line separation between them. The right sinuses involvement is commoner than the left. These results agree with some previous studies.<sup>12,14</sup> Preseptal cellulitis was the most infection found in our series in 53.3% of cases, and treated on outpatient basis. Radovani et al<sup>6</sup> reported that the most commonly occurring complication is palpebral inflammatory oedema (preseptal cellulitis), it encountered more often in children. The upper eyelid becomes swollen and hyperaemic due to a blockage of vein drainage from an ethmoid sinusitis. Chemosis and proptosis are absent, both of which usually indicated postseptal infection. Vision remains unaffected and the eyeball moves in all directions.<sup>4,6</sup> Another study reported that periorbital abscess was the most commonly complications secondary to rhinosinusitis was occurred in 58% of cases.<sup>14</sup> This finding against our result. Periorbital cellulitis was found in 20% of our series, all of them treated medically with success rate 100%.

This result is similar to previous studies which reported that orbital cellulitis was occurred in 22% of cases and treated medically.<sup>11,14</sup>

Unfortunately, when a patient presents with periorbital cellulitis it is not easy to defined which stage the disease has reached on clinical signs alone and it must be remembered that half of cases with intracranial complications present

with periorbital cellulitis.<sup>15</sup> Orbital cellulitis occurs when infection spreads posterior to the orbital septum and can lead to abscess if not adequately treated. Radiological examination is required when orbital abscess is suspected.<sup>8,16</sup> Periorbital abscess is collection of pus between the lamina papyracea and the orbital periosteum. Proptosis, chemosis, and limited movement of the eye ball are present. Radovani et al<sup>6</sup> reported that all cases in this series were underwent surgical intervention under the protection of combined antibiotherapy. The appearance of proptosis, as well as chemosis, indicated that the spread an inflammatory process in the anterior part of the orbit. In our study, there was 20 % of cases presented with subperiosteal abscess, 2 cases was presented with early and small subperiosteal abscess those treated medically, by I.V. antibiotics and locally decongestant with good results whereas 4 cases treated by combined modalities, surgical intervention through nasal endoscopy under covering of antibiotics. Previous studies reported that all cases with subperiosteal abscess were treated by surgical drainage.<sup>8,9,17</sup> There have been literatures<sup>18,19</sup> reported on the role of vigorous medical therapy alone for post-septal involvement in selected cases -the selected criteria proposed by Oxford et al<sup>19</sup> includes normal vision, absence of ophthalmoplegia, intraocular pressure < 20 mmHg, proptosis of 5 mm or less and an abscess width of 4 mm or less on CT scan. In our study, there was one case presented with intraorbital abscess, and other case with cavernous sinus thrombosis presented with bilateral orbital swelling, these two cases were treated by combined modalities therapy, surgical and medical. In spite of vigorous medical treatment and surgical drainage by external approach,

intraorbital abscess was ended with loss of vision and cavernous sinus thrombosis patient expired after one week of admission to the hospital.

An accumulation of pus begins in the retro-orbital adipose tissue. This is a more severe complication because it makes up the precedent for the thrombosis of the cavernous sinus, or passing of the infection through nervous route into the intracranial space.<sup>6</sup> One patient was presented with fistula of the right ethmoid sinus, the parents refused surgical intervention as first line of the treatment at the time of presentation, later on this patient was underwent to external ethmoidectomy with good prognosis. One of the main clinical indications of surgery is imaging that demonstrates an abscess or potential compromise of critical structures, such as the eye or brain.<sup>20</sup> Signs that indicate abscess formation include, decrease in vision, proptosis, ophthalmoplegia and pain associated with eye movement.<sup>11</sup>

Orbital complications of sinogenic origin should always be treated as emergency and treated aggressively as it poses life-threatening intracranial complication and blindness. Recognition, correct diagnosis, and appropriate treatment is necessary to avoid orbital complications of sinogenic origin.<sup>21,22</sup> Rhinosinusitis is the most common cause of orbital infection in the children. Orbital complications secondary to rhinosinusitis can result in permanent blindness or death if not treated promptly and appropriately.<sup>23,24</sup>

The outcomes of treatment depended on the number and types of complications and general condition of the patient.<sup>24</sup> Early presentation, effective antibiotics and co-management with other specialist improved outcome. There is need for health education to avoid over-counter

drugs, treat predisposing problem such as sinusitis and dental infection.<sup>25,26</sup>

### Conclusion

Orbital complications secondary to rhinosinusitis are the most common in the children. Orbital Complications can result in permanent blindness or death if not treated promptly and appropriately. The early recognition and aggressive treatment of orbital complications are the mainstay to reduction of unwanted complications and morbidity. The majority of patient were treated medically with good responding, while surgical procedure is done for small group of the patients.

### References

1. Hermann BW, Forsen JW. Simultaneous intracranial and orbital complications of acute rhinosinusitis. *Int J Pediatr Otolaryngol* 2004; 68: 619-62
2. Chandler JR, Langebrunner DR, Stevens ER. The pathogenesis of orbital complications in acute sinusitis. *Laryngoscope* 1970;80:1414-28
3. Awaorgu OGB, Awoben FJ, Onakoya PA, Awoben AA. Orbital cellulitis complicating sinusitis: A 15 years review. *Nigeria J Surg Research* 2004;6 (1): 14-16
4. Moloney JR, Badham NJ, McRae A. The acute preseptal (peri-orbital) cellulitis, subperiosteal abscess and orbital cellulitis due to sinusitis. *J Laryngol Otol* 1987;101: (suppl12): 1-14
5. Swift AC, and Charlton G. Sinusitis and the acute orbit in children. *J Laryngol Otol* 1990; 104:213-16
6. Radovani P, Vasili D, Xhelili M, Dervishi J. Orbital complications of sinusitis. *Balkan Med J* 2013;30:151-4
7. Ali A, Kurien M, Mathews SS, Mathew J. Complications of acute infective rhinosinusitis experience from a developing country. *Singapore Med J* 2005; 46:540-4
8. Suhaili DN, Goh BS, Gedeh BS. A ten years retrospective review of orbital

- complications Secondary to acute sinusitis in children. *Med J Malaysia* 2010;65 (1):49-52
9. Siede K V, Kemer A, Betz CS, Tschiesner U, Berghaus A, Leunig A. Management of orbital complications due to rhinosinusitis. *Eur Arch Otorhinolaryngol* 2010;267:1881-6
  10. William SR, Carruth JAS. Orbital infection secondary to sinusitis in children: diagnosis and management. *Clin Otolaryngol* 1992;17:550-557
  10. AL-Madani MV, Khatatbeh AF, Rawashdeh RS, AL-Khoum NF, Shawagefeh NR. The prevalence of orbital complications among children and adults with acute rhinosinusitis. *Braz J Otorhinolaryngol* 2013;79.
  11. Pena MT, Preciado D, Orestes M, Choi S. Orbital complications of acute sinusitis change in the post-pneumococcal vaccine Era. *JAMA Otolaryngol Head Neck Surg* 2013;139(3):223-227
  12. Mortimore S, Wormald PJ. The Goote Schuur hospital classification of the orbital complications of sinusitis. *J Laryngol Otol* 1997; 111:719-723.
  13. Peter A, Mohammed D I. Results of orbital cellulitis treatment. *J Natural Scien Rerarch* 2012;2(5): 91-5
  14. Jones NS, Walker JL, Bassi S, Jones T, Punt J. The intracranial complications of rhinosinusitis: can they be prevented? *Laryngoscope* 2002;112:59-63
  15. Bolinaga U, Petez N, Labato R, Martinez-Seijas P, Sumtier Perez E. Abscess of the orbit as a complication of acute sinusitis: surgical approach. *Eur Arch of otorhinolaryngol* 2009;266:(A368):1037-8
  16. El-Silimi O. The place of endonasal endoscopy in the treatment of orbital cellulitis. *Rhinology* 1995;33:93-6.
  17. Green MF, Polard ZF. Medical treatment of pediatric subperiosteal orbital abscess. *Jaapos* 1998;2:351-55.
  18. Oxford LE, McClay J. Medical and surgical management of subperiosteal orbital abscess secondary to acute sinusitis in children. *Int J Pediatr Otorhinolaryngol* 2006;70:1853-6.
  19. Dhangra BA, Linder L, Preciado DA, Taylormoore J, Shah RK. Orbital preorbital infections Anational perspective. *Arch Otolaryngol Head Neck Surg* 2011;137 (8):769-73.
  20. Chavan SS, Deshmukh DS, Pawar VG, Kirpan VG, Khobragade SW, Sarvade k, et al. Orbital Complications of sinogenic origin: A case study of 20 patients. *Word articles in ear, nose, and throat* 2010; 3. [www.waent/archives/2010/vol3-2/10.html](http://www.waent/archives/2010/vol3-2/10.html).
  21. Rashed F, Canon A, Heaton PA, Pual SP. Diagnosis, management and treatment orbital and periorbital cellulitis in children. *Emerg Nurse* 2016;24(1) :30-5.
  22. Singh CSK, James SCE, Sabarigirish CK, Sawami WH, Sood MT. Bilateral orbital complications of pediatric rhinosinusitis. *MJAFI* 2014; 70:68-72.
  23. Chaiyate S, Foonant S, Navacharoen N, Roongrotwattanasiri K, Pongsakom T, Patumanond J. Complications of sinusitis in a tertiary Care Hospital: types, patients
  24. Characteristics and outcomes. *Int J Otolaryngol* 2015
  25. Wan Y, Shi G, Wang H. Treatment of orbital complication following acute rhinosinusitis in children. *Balkan Med J* 2016;33 (4) :401-6.
  26. Adeoti CO, Adejumo OO, Isawumi MA, Kolawole OO, Haastrup AA. Orbital cellulitis in a tertiary institution. *Niger J Ophthalmol* 2017;25: 141-5



# Al-Razi University Journal for Medical Sciences

صنعاء - شارع الرباط - خلف البنك اليمني للإنشاء والتعمير

فاكس : 406760

تلفون : 216923

[www.alraziuni.edu.ye](http://www.alraziuni.edu.ye)

[info.alraziuni.edu.ye](mailto:info.alraziuni.edu.ye)