

# Al-Razi University Journal for Medical Sciences




## RUJMS

Print ISSN No. 2616-6143

Online ISSN No. 2708-0870

Volume (4) Issue (2) DEC 2020



مرخصة من وزارة التعليم العالي والبحث العلمي  
بقرار وزاري رقم ( 57 ) لسنة 2018م

### RUJMS

Published by Al-Razi University

Biannual Referred Journal

All Rights Reserved for Al-Razi University

Editor In Chief	Nationality	Degree
Dr. Nabil Ahmed Al-Rabeei	Yemen	Professor
Editor Manager	Nationality	Degree
Dr. Mohammed S.A. Al-Awar	Yemen	Associate Professor

### Editorial Board Members

No	Editorial Board Members	Nationality	Degree
1.	Dr. Abdulsalam. M. Dallak	Yemen	Professor
2.	Dr. Abduljalil D. Ghaleb	Yemen	Professor
3.	Dr. Mohammed A. Haidar	Yemen	Professor
4.	Dr. Mahfouth A. Bamashmus	Yemen	Professor
5.	Dr. Mohammed Aissa	Yemen	Professor
6.	Dr. Ahmed Al-Sobati	Yemen	Professor
7.	Dr. Abdulhameed Al-Thifani	Yemen	Associate Professor
8.	Dr. Nouradden Al-Jaber	Yemen	Associate Professor
9.	Dr. Shatha Hassan Yassin	Yemen	Associate Professor
10.	Dr. Sadeq Hassan Al-Sheraji	Yemen	Associate Professor
11.	Dr. Ahmed Ali Abdulateef	Yemen	Associate Professor
12.	Dr. Abdulmajid Alssaifi	Yemen	Associate Professor

### Advisory Board

No	Advisory Board	Nationality	Degree
1.	Dr. El Houcin Boidida	Morocco	Associate Professor
2.	Dr. Yahia Cherrah	Morocco	Associate Professor
3.	Dr. Abdulaziz Benjouad	Morocco	Associate Professor
4.	Dr. Abdellah Akil	USA	Associate Professor
5.	Dr. Katim Alaoui	Morocco	Associate Professor
6.	Dr. Arvinder Bahala	India	Associate Professor
7.	Dr. David Tasala	USA	Associate Professor

**Copyright of articles published in the RUJMS belong to the University of Al-Razi unless the work is subject to copyright.**

**Address: Al-Razi University - College of Medical Sciences**

**Telefax: +9671406760 P.O. Box:1152 Sana'a – Yemen**

**Website: <http://alraziuni.edu.ye/rujms/>**

**Designed by Eng. Osama Al-Moaina  
Ossamah245@yahoo.com**



## Toxic Effect of Methidathion and Chlorpyrifos on Liver of Male Rabbits

\*<sup>1</sup>Adel A.A Muaqeb, <sup>2</sup>Mohammed S. Al-Awar, <sup>1</sup>Ebtisam Y. Shekoo

<sup>1</sup>Department of Biology, Faculty of Education, Aden University. <sup>2</sup>Department of Medical Laboratory, Collage of Medical Science, Al-Razi University, Yemen

\*Corresponding author: Adel A.A Muaqeb, Department of Biology, Faculty of Education and Science, Aden University. Email: dradelmuaqeb@gmail.com

### Abstract:

The irrational use of organophosphorus insecticides in Yemen play a crucial role in the occurrence of many diseases affecting plants, animals, and man. Methidathion (MD) and Chlorpyrifos (CF) are two of widely used organophosphorus insecticides in agriculture. The purpose of the present study was to investigate the in vivo effect of (MD) and (CF) on lipid peroxidation (LPO), antioxidant defense system in the liver cells and liver function in male rabbits after orally administration a single dose of 1/8 of LD50 of (MD) and (CF) period of 20 days. Thirty healthy male rabbits weighting 1400-1600g., were divided into 3 groups with 10 animals in each, first group served as control animals, they received 3ml. of corn oil, while animals in second and third groups received 1/8 of LD50 of (MD) and (CF) respectively. The level of alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), lactate dehydrogenase (LDH), and gamma glutamyl transferase ( $\gamma$ -GT) in serum were analyzed, the level of (LPO), and methylglyoxal (MG), in liver homogenate were also studied. Results showed a significant ( $P < 0.01$ ) increase in all studied parameters in animals treated with (MD) and (CF) as compared to control animals. Our results indicated to the high toxicity of (MD) and (CF).

**Key words:** Methidathion, Chlorpyrifos, Lipid peroxidation, Liver. Introduction

DOI: <https://doi.org/10.51610/rujms4.2.2020.58>

### Introduction

The pollution of the environment plays a crucial role in the occurrence of many diseases affecting plants, animals and man. One of the main factors causing pollution of the environment is the irrational use of organophosphorus insecticides<sup>1</sup>. Many alterations have been observed in organs of animals due to the organophosphorus insecticides. The organophosphorus insecticides have been implicated as playing a certain role in etiology of tumors<sup>6</sup>, blood diseases<sup>10</sup>, liver injury<sup>4,12,21</sup>, heart and vascular muscles atrophy<sup>26,27</sup>, digestive system

diseases<sup>6</sup>, renal disorders [5 and 6], and metabolism disorders<sup>5</sup>. Methidathion (MD) and Chlorpyrifos (CF) are an organophosphorus insecticides with anticholinesterase mode of action<sup>13,26</sup>, (MD) and (CF) are two of the most widely and irrationally used insecticides in agriculture in Yemen, especially in khat cultivation<sup>1</sup>. It has been found that the subchronic orally administration of (CF) significantly increased the oxidative stress in rats erythrocytes caused their haemolysis<sup>13</sup>. The orally administration of (CF) resulted in significant increase in the level of thiobarbituric acid reactive substances (TBARS), and significant decrease in

superoxide dismutase and catalase in rats lung tissues<sup>18</sup>. White blood cells, red blood cells, hemoglobin, and total protein significantly decreased, while creatinine and uric acid significantly increased in blood of mice orally treated with (CF)<sup>5</sup>. The orally administration of (MD) increased the oxidative stress and decreased the antioxidant enzymes activities in rats erythrocytes<sup>3</sup>. Histopathological change like infiltration with mononuclear cells at parenchymal tissue, sinusoidal dilatation, focal necrotic areas, granular degeneration and picnotic nuclei in the hepatocytes were observed in rats liver after orally administration of (MD)<sup>12</sup>. The level of cardiac troponin in serum and MDA in cardiac tissues significantly increased in rats after orally administration of (MD), this indicates to the diffuse loss of striation and myocytolysis of the cardiomyocytes caused by (MD)<sup>26</sup>.

The purpose of the present study was to investigate the *in vivo* effect of (MD) and (CF) on lipid peroxidation (LPO) and antioxidant defense system in the liver cells and on liver function in male rabbits.

## Materials and Methods

### Animal's treatment and blood collection:

Thirty healthy male rabbits local breed (1400-1600g) were divided into 2 treated groups and control, as follows:

**Control group:** 10 animals treated with a single daily dose of 3ml.corn oil orally period of 20 days.

**(MD) group:** 10 animals treated with a single daily dose of 1/8 of LD<sub>50</sub> of (MD) (5mg/kg) in 3ml. corn oil orally period of 20 days.

**(CF) group:** 10 animals treated with a single daily dose of 1/8 of LD<sub>50</sub> of (CF) (12mg/kg) in 3ml. corn oil orally period of 20 days.

All animals were maintained in standard environmental conditions and kept a standard commercial diet with water *ad libitum*.

The experiment was administrated in the Animal Physiology Laboratory, Department of Biology, Faculty of Sciences and Education,

and Central Researches Laboratory Aden University.

After 20 days the animals were fasted over night for 12hrs, then they were sacrificed, the blood was immediately collected and centrifuged, serum was discarded and kept at -21 °C for the biochemical analyze. Livers were removed immediately after death, perfused with normal saline containing heparin, homogenized with phosphate buffer saline (pH 7.2) using homogenizer, centrifuged at 3000g for 30 min. The supernatant was removed and stored at -20°C.

## Analysis

### Alanine-aminotransferase and Aspartate-aminotransferase Assay

The level of ALT and AST in serum was determined according to the procedure described by Gehringer *et al.*,<sup>11</sup>.

### Gamma glutamyl transferase Assay

Determination of  $\gamma$ -GT activity in serum was based on the method of Szasz<sup>22</sup>.

### Lactate dehydrogenase Assay

Serum LDH was measured according to Doyle and Griffiths<sup>9</sup>.

### Alkaline phosphatase Assay

ALP was determined using a colorimetric method as described by Kind and King<sup>19</sup>.

### Lipid peroxidation Assay

The lipid peroxidation (LPO) level in hepatocytes homogenate was measured via the measuring of malondialdehyde MDA according to method described by Husseinzadeh *et al.*,<sup>15</sup>.

### Methylglyoxal Assay

The MG was determined in the liver homogenate according to the method described by Ratliff *et al.*,<sup>20</sup>.

### Statistical analysis

The statistical analysis was performed by SPSS; continuous data are expressed as mean  $\pm$  S.E. Data were compared using one – way

ANOVA. P value <0.01 was considered to be statistically significant.

## Results

Table1.The level of enzymes in serum after 20 days of orally administration of (MD) and (CF) in dose 5mg/kg and 12mg/kg respectively.

Enzymes	Treatments		
	Control	MD	CF
ALT IU/L	41.35 $\pm$ 1.32	141.25 $\pm$ 4.43	120.10 $\pm$ 2.11
AST IU/L	32.20 $\pm$ 1.65	185.10 $\pm$ 3.85	161.20 $\pm$ 5.12
ALP IU/L	59.65 $\pm$ 2.73	139.60 $\pm$ 6.34	105.82 $\pm$ 3.92
LDH IU/L	210.25 $\pm$ 4.12	360.34 $\pm$ 4.11	302.61 $\pm$ 6.54
$\gamma$ -GT IU/L	6.13 $\pm$ 0.12	36.11 $\pm$ 1.14	22.16 $\pm$ 1.09

Values are expressed as mean of 10 animals $\pm$ S.E, P<0.01 vs. control.

Table 2.The level of MDA and MG in liver homogenate after 20 days of orally administration of (MD) and (CF) in dose 5mg/kg and 12mg/kg respectively.

Parameters	Treatments		
	Control	MD	CF
MDA $\mu$ M	0.03 $\pm$ 0.01	18.23 $\pm$ 2.11	13.21 $\pm$ 3.43
MG $\mu$ M	1.12 $\pm$ 0.33	59.18 $\pm$ 4.87	41.02 $\pm$ 4.86

Values are expressed as mean of 10 animals $\pm$ S.E, P<0.01 vs. control.

Data in table1 show that, the administration 1/8 of LD<sub>50</sub> (MD) and (CF) leads to the critical damage in liver function. The results in table1 show high significant (P<0.01) increase in the level of aminotransferases ALT and AST, as well as in ALP,LDH and  $\gamma$ -GT, in the serum of animals treated with (MD) and (CF) as compared to control animals, this increase was higher in the serum of animals treated with (MD).

Data in table2 show high significant (P<0.01) increase in the level of MDA a product of lipid peroxidation (LPO) and in MG a toxic substance in the liver homogenate of treated with (MD) and (CF) animals as compared to control animals, this increase was higher in the serum of animals treated with (MD).

## Discussion

According to the literatures the increase in AST and ALT indicates to acute liver diseases, cirrhosis liver and toxic liver cells necrosis<sup>7,14</sup>, while the increase in LDH indicates to acute hepatitis, renal tubular necrosis and acute myocardial infraction<sup>7</sup>. The increase in ALP is due to the lesions of liver, ALP also increased in acute liver diseases<sup>7</sup>. The  $\gamma$ -GT increases in the most of hepatobiliary<sup>7</sup>. These facts suggested that the exposure to (MD) and (CF) in our experiment led to serious damage in liver cells.

One important consequence of exposure to (MD) and (CF) is the excessive of free radicals

production, which attack many organic molecules in cells membrane including polyunsaturated fatty acid leading to increase in LPO<sup>8</sup>. Several studies reported that reactive oxygen species (ROS) initiate LPO through the action of hydroxyl radicals<sup>16</sup>. Exposure to the toxic compounds has been linked with increase of ROS production not only in domestic and wildlife animals but also in human<sup>25</sup>. Formation of reactive oxygen species (ROS) and oxidative stress is associated with development of many pathological statuses<sup>24</sup>. Oxidative stress may occur either due to the decrease of cellular antioxidant level, or due to the over production of ROS<sup>16</sup>.

Our results clearly showed that (MD) and (CF) in the negligible dose have the ability to free radicals formation, decrease of cellular antioxidant level, therefore initiate of the LPO which leading to damage of cells membrane and death of cells<sup>24</sup>. Our results are close to that found in previous studies<sup>3, 4, 12</sup>.

Methylglyoxal (MG), (also called Pyruvaldehyde), formed as a side product of dihydroxyacetone phosphate in glycolysis, MG must converted into lactic acid and/or pyruvic acid in presence of glyoxalase I<sup>7</sup>. MG is considering being toxic for mammals, the MG cytotoxicity is due to it accumulation in cells<sup>17</sup>. The animals exposure to (MD) and (CF) in our experiment showed increase in the MG level in liver cells homogenate. The accumulation of MG is due to the inhibition of glyoxalase I<sup>17, 20</sup>, the accumulation of MG in liver caused depletion of antioxidant defense system found in liver, leading to significant generation of free radicals which might in further strengthen the damage and affect the hepatocytes function<sup>2, 17, 23</sup>. Our results clearly showed that the exposure to the organophosphorus insecticides (MD) and (CF) in negligible dose inhibited the glyoxalase I, therefore, accumulation of MG in liver cells, that leads to depletion the antioxidant defense system, caused significant increase in free radicals generation; which leads to increase in LPO, thus damage of hepatocytes function, and so increase in the level of enzymes concerning liver dysfunction. The high toxicity

of organophosphorus insecticides may due to their ability to free radicals formation. The free radicals have been implicated as playing a major role in induce oxidative stress, therefore, the etiology of many diseases and alterations<sup>24</sup>.

## References

1. Al-Haj M. Awadh N.A. and Ali A.A. (2005). Survey of pesticides used in Qat cultivation in Dhale and Yafe and their adverse effects. *J. Nat. Appl. Sci.* 9 (1):103-110.
2. Al-Jassabi S. (2004). Effect of methylglyoxal on antioxidant enzymes of the liver and spleen of mice. *J. Biol. Sci.* 4 (5):605-608.
3. Altuntas I. Delibas N. and Sutcu R. (2002). The effect of organophosphorus insecticide methidathion on lipid peroxidation and antioxidant enzymes in rat erythrocytes: role of vitamins E and C. *Hum. Exp. Toxicol.* 21 (12):681-685.
4. Altuntas I. Delibas N. Demirci M. Kiline I. and Tamer N. (2002). The effect of methidathion on lipid peroxidation and some liver enzymes: role of vitamins E and C. *J. Toxicol.* 76(8):470-473.
5. Ambali S. Akanbi D. Igbokwe N. Shittu M. Kawu M. and Ayo J. (2007). Evaluation of subchronic chlorpyrifos poisoning on hematological and serum biochemical changes in mice and protective effect of vitamin C. *J. Toxicol. Sci.* 32 (2):111-120.
6. Bhatnagar J.A. 1980. Effect of pesticide stress among pesticide factory workers in Delhi. *Pub. HIM.* 74:380.
7. Chatterjea M.N. and Shinde R. (2005). Text Book of Medical Biochemistry. 6th ed. Jaypee Broth. New-Delhi. pp641-644.
8. Denisov E.T. Denisova T.G. and Pokidova T.S. (2003). Hand book of free radical initiators. John Wiley & Sons Inc. Hoboken, New-Jersey, USA, pp. 22-57.
9. Doyle A. and Griffiths J.B. (1988). Cell and tissue culture: laboratory procedures in biotechnology. John Wiley & Sons, Baffins Lane Chichester, West Sussex, England pp.71-74.
10. Ebram R.A. and Sahati M.A. (2009). The organophosphorus pesticides, pollution and toxicity. 2nd ed. Puna pub. Inc. Puna,

- India. Pp.122-124
11. Gehringer M.M. Downs K.S. Downing T.G. Naude R.G. and Shephard E.G. (2003). An investigation into the effect of selenium supplementation on microcystin hepatotoxicity. *Toxicol.*41:1-8.
  12. Gokalp O. Gulle K. Sulako O. Cicek E. Altuntas I.(2003).The effect of methidathion on liver : role of vitamins E and C. *Toxicol.Ind.health* 19(2-6):63-67.
  13. Gultekin F. Delibas N. Yasar S.and Kilinc I.(2001).In vivo change in antioxidant system and protective role of melatonin and a combination of vitamin C and vitamin E on oxidative damage in erythrocytes induced by chlorpyrifos in rats. *Arch.Toxicol.*75 (2):88-96.
  14. Guyton C. and Hall E.(2006).Text Book of Medical Physiology.11th ed.Elsiver Inc.Pennsylvania, USA.pp.511-512.
  15. Hosseinzadeh H. Paravardeh S. Asl M.N. Sadeghnia H.R. and Ziaee T.(2007). Effect of thymoquinone and *Negella sativa* seeds oil on lipid peroxidation level during global cerebral ischemia – reperfusion injury in rats' hippocampus. *Phytomed.*14:621-271.
  16. Joshi G. Sultana R.Tangpong J. Cole M.P. Clair D. Vore M. Estus S. and Butterfield D.A.(2005).Free radical mediated oxidative stress and toxic side effects in brain induced by the anticancer drug adriamycin : insight chemobrain.*Free Rad.Res.*39(11):1147-1154.
  17. Kalapos M.P.(2008).The tandem of free radicals and methylglyoxal. *Chemico.Biol.Inter.*171 (3):251-271.
  18. Karaoz E.Gultekin F. Akdogan M. Oncu M. and Gokcimen A.(2002).Protictive role of melatonin and a combination of vitamins C and E on lung toxicity induced by chlorpyrifos in rats.*Exp.Toxicol.Pathol.*54(2):97-108.
  19. Kind P.R. and King E.G.(1954).Estimation of plasma phosphate by determination of hydrolysed phenol with amino-antipyrine.*J.Clin.Path.*7:56-63.
  20. Ratliff D.M.Vander Jarget D.J. Eaton R.P. and Vander Jarget D.L.(1996).Increase level of methylglyoxal-metabolizing enzymes in mononuclear and polymorphonuclear cells from insulin-dependent diabetic patients with diabetic complications: aldose reductase glyoxalaseI and glyoxalaseII. *J.Clin.Endocrinol.Metabol.*81 (2):488-492.
  21. Sutcu R. Altuntas I. Yildirim B. Karahan N. Demirin H. and Delibas N.(2006).The effect of subchronic methidathion toxicity on rats liver: role of antioxidant vitamins C and E. *Cell Biol.Toxicol.*22(3):221-227.
  22. Szasz G. (1974).Methods of enzymatic analysis. *Natural. Toxins* 2:714-715.
  23. Thornalley P.G.(2003).Glycolase I – structure, function and critical role in the enzymatic defense against glycation. *Biochem.Soc.Trans.* 31(6):1343-1348.
  24. Valko M. Leibfritz D.Moncol J. Cronin M. Mazur M. and Telser J. (2007). Free radicals and antioxidants in normal physiological functions and human diseases. *Int.J.Biochem.Cell Biol.*39 (1):44-84.
  25. Weng D. Lu Y.Wei Y. LIU Y. and Shen P.(2007).The role of reactive oxygen species (ROS) in microcystin -LR-induced hepatocytes apoptosis and liver injury in mice.*Toxicol.*232:15-23.
  26. Yavuz T. Altuntas I. Delibas N. Yildirim B. Candir O. Cora A. Karahan N. Ibrisim E. and Kutsal A. (2004).Cardiotoxicity in rats induced by methidathion and ameliorating effect of vitamins E and C. *Hum.Exp.Toxicol.*23(7):323-329.
  27. Yavuz T. Altuntas I. Delibas N. Yildirim B. Candir O. Cora A. Karahan N. Ibrisim E. and Kutsal A. (2004). Vascular wall damage in rats' damage induced by methidathion and ameliorating effect vitamins C and E. *Arch.Toxicol.*78 (11):655-65.



## Author Guidelines



RUJMS

# Al-Razi University Journal of Medical Sciences



### Manuscript Preparation

The manuscripts should be in word document (Microsoft Word 97, 2003, 2007). Manuscripts, well-written in English.

- **Fonts:** To distinguish different parts of the article, it is recommended to use the font single space, Times New Roman, 14 bold for the title, size 10 for the abstract, size 12 for the body, size 12 **bold** for headings and size 11 for references.

### Paper Title

Choose a title that is long enough to cover the main points and by considering the importance of shortness. Only the first letter of the first word of each title.

### Authors & Affiliations

- Author(s): full name of all authors should be mentioned: [First Name] [Middle Name] [Last Name]
- Affiliation: Author's affiliation contains only [name of department], [name of institution]
- Corresponding author: Their address, telephone and email should be written.

### Abstract and Keywords

- **Abstract:** Required for original articles, review articles, case reports
  - An abstract of 250 words should be provided.
  - Abstract of should be structured under the following headings: Background, Aim, Methods, Results and Conclusions.
- **Keywords:** A maximum of 5 keywords must be provided for indexing purposes.

### Text

*For original articles* the manuscript divided into the following headings: introduction, Aim, Methods, Results, Discussion and conclusion.

- **Introduction:** This section should be succinct, with no subheadings.
- **Aim:** This should be follows introduction.
- **Subjects and Methods:** This part should contain sufficient detail so that all procedures can be repeated
- **Results:** This section may each be divided by subheadings or may be combined.
- **Discussion:** This section may each be divided by subheadings or may be combined.
- **Conclusion:** This should clearly explain the main conclusions of the work highlighting its importance and relevance.

- **Recommendations:** should be apply
- **Acknowledgements:** All acknowledgments should be included at the very end of the paper before the references and may include supporting grants, presentations, and so forth.

---

## References

---

This Journal accepts references according to a style based on Vancouver.

References: 25 – 30 should be provided.

### *In - text references*

---

In the Vancouver system, the only indication required in the text of a paper is a number, allocated in ascending sequence, and presented in the text either in brackets, some or in superscript.

- **Citing one piece of work,** For example:  
Recent research<sup>1</sup> indicates that the number of duplicate papers being published.
- **Citing more than one piece of work at the same time,** example:  
Multiple clinical trials<sup>4-6,9</sup> show.....
- **Citing the author's name in your text,** example:  
Al-Rabeei<sup>8</sup> has argued that.

### *In reference list*

---

#### **Book: print**

- **Standard Format for Books:**  
Author Surname, Initials. Title: subtitle. Edition (if not the first). Place of publication: Publisher; Year.

#### **Examples:**

- **Book with One Author or Editor, example:**  
Snell RS. Clinical anatomy by regions. 9th ed. Philadelphia: Lippincott Williams & Wilkins; 2012.
- **Two-Six Authors/Editors, example:**  
Simons NE, Menzies B, Matthews M. A Short Course in Soil and Rock Slope Engineering. London: Thomas Telford Publishing; 2001.
- **More than Six Authors/Editors, example:**  
Fauci AS, Braunwald E, Kasper DL, Hauser SL, Longo DL, Jameson JL, et al., editors. Harrison's principles of internal medicine. 17<sup>th</sup> ed. New York: McGraw Hill; 2008.

#### **Articles in Journals: print**

- **Standard Format for Journal Articles:**  
Author Surname, Initials. Title of article. Title of journal. Date of Publication: Volume Number (Issue Number): Page Numbers.

#### **Examples:**

- **Journal article with one author**  
Al-Rabeei N. Analyze of care accidents in Sana'a City. *Indian J of Pub. H.* 2009;74(4):12-15.
- **Journal article with Two-Six Authors, example:**  
Al-Rabeei N, Dallak A, Al-Barakani M. Complementary feeding of infants and young children in Sana'a City, Yemen. *EMJ.* 2014;12(2):13-19.

- ***Journal article with more than six authors***

Dallak A, Al-Rabeei N, Al-Barakani M, Al-Thifani A, Al-Rabeei J, Hazam S, et al. Prevalence of malnutrition among infants in Sana'a city, Yemen. *Ulutas Med J.*2015;1(1):1-6.

### **Tables/Figures**

---

- Tables / Figures: should be 4–5
- Each Table should have borders with normal style without any colour row or column.



# Al-Razi University Journal for Medical Sciences

صنعاء - شارع الرباط - خلف البنك اليمني للإنشاء والتعمير

فاكس : 406760

تلفون : 216923

[www.alraziuni.edu.ye](http://www.alraziuni.edu.ye)

[info.alraziuni.edu.ye](mailto:info.alraziuni.edu.ye)